
The geodesic climate model of the (near) future







David Randall, CSU Todd Ringler, CSU Ross Heikes, CSU Cara-Lyn Lappen, CSU Wayne Schubert, CSU Akio Arakawa, UCLA Celal Konor, UCLA Albert Semtner, Jr., NPGS Scott Fulton, Clarkson University





What's up

We promised to develop and deliver to the Climate Change Prediction Program a *geodesic* coupled ocean-atmosphere-land surface model designed for climate applications.

The atmosphere and ocean will use hybrid quasi-Lagrangian vertical coordinates.

The computational architecture of the model will be designed for use on emerging new computer architectures.

A key goal of our project is to train young climate modeling researchers in all aspects of the climate modeling enterprise.






The Climate system Changes of Solar radiation radiation H₂O, N₂, O₂, Clouds CO₂, O₃, etc., **Atmosphere-land** aerosols coupling Atmosphere-ice coupling Wind stress exchange Precipitation Heat Evaporation 🌡 Ice-ocean Atmosphere-ocean coupling J coupling Changes of atmospheric compositor Changes of land features Changes of ocean basin orography, vegetation, albedo, etc shape, salinity, etc.





Computational structure



The flux coupler is "the boss."





Framework

We started with a blank piece of paper.


- All model components use geodesic grids.
- Atmosphere and ocean both use quasi-Lagrangian vertical coordinates.
- Parallel flux coupler takes advantage of similar grids.
- Physical parameterizations are tailored to work with the numerical framework.


All components are home-grown.





Geodesic grid(s)





All cells are nearly the same size.

The grid is quasi-isotropic.

Advective CFL is not much of an issue.







Generating the grid


Icosahedron

Bisect each edge and connect the dots


Pop out onto the unit sphere








.... and so on, until we reach our target resolution.











Quasi-Lagrangian Vertical Coordinates



sigma

Lower boundary is a coordinate surface.

Mass flows freely across layer edges.




theta

Lower boundary is not a coordinate surface.

Mass stays within layers except due to heating.

Cannot represent static instability.



zeta

Lower boundary is a coordinate surface.


Becomes theta smoothly and naturally.

Automatically reverts to sigma in statically unstable layers.






Embedded boundary layer








Mass exchanges across the PBL top








Clouds and stuff







Ocean and sea-ice models




In the sea ice model, transport is represented with a remapping algorithm.

The same approach will be tested in the ocean model.





Land-surface model







Flux coupler

The atmosphere is represented on a coarse geodesic grid, while the ocean and land-surface are represented on a finer geodesic grid.

Exchanges of energy, mass, and momentum are computed on the fine grid.

The atmospheric state is "interpolated down," and the fluxes are "averaged up," using methods developed by Phil Jones of LANL.



We are not interested in coupling all possible grids to all possible grids.


We are taking advantage of the compatibility of our surface and atmosphere grids.





Parallelization with MPI

Two-dimensional data decomposition works like this:



40 panels



160 panels

Each color corresponds to a processor. In this simplified example there are only four processors.





Animation





Status

• The generalized coordinate is being tested in the atmosphere model.

The AGCM is nearing completion.

• The sea ice model and land-surface model are well along.

These are closely related to the corresponding components of the CCSM.

• The ocean model is lagging somewhat but we are pushing on this now.

Major progress is expected this year.



